

Computer processing of lung scintigraphic images

N.I. Glumov^{1,2}, A.V. Kapishnikov²

¹Image Processing Systems Institute of RAS

²Samara State Medical University

Abstract

The authors have developed the methods of computer processing of scintigrams allowing to objectify the state of capillary blood flow in lungs by a set of computer parameters: the spectral characteristics of activity nonuniformity and the measure of perfusion deficiency. The results of clinical studies, confirming high efficiency of the developed algorithms, are presented.

Keywords: lung scintigraphic, computer processing, image, capillary blood, perfusion deficiency, nonuniformity.

Citation: Glumov NI, Kapishnikov AV. Computer processing of lung scintigraphic images. Computer Optics 2003; 25: 158-164.

[Access full text \(in Russian\)](#)

References

- [1] Kosarev VV, Kapishnikov AV. Radionuclide diagnostics of regional perfusion disorders in lungs in pulmonary dust disease [In Russian]. Proceedings of the 7th National Congress of Respiratory Diseases 1997: 306.
- [2] Soifer VA, ed. Methods of computer image processing. Moscow: "Fizmatlit" Publisher; 2001.
- [3] Koroluk IP, Kapichnikov AV. Evaluation of alveolar permeability, lung ventilation and perfusion in pneumoconiosis. Eur J Nucl Med Mol Imaging 1998; 25(9): 1186.
- [4] Krause M, Kummer G, Mahrlein W. Zusammenhang zwischen den pulmonalarterien mitteldruck enunterruehe- und belastungsbedingungen in abhangigkeit vom silikosestadium. Z Erkr Atmungsorgane 1985; 164(3): 288-291.
- [5] Jandová R, Widimský J, Eisler L, Navrátil M. Long-term prognosis of pulmonary hypertension in silicosis. Cor Vasa 1980; 22(4): 221-237.
- [6] Kitabatake A, Inoue M, Asao M, et al. Noninvasive evaluation of pulmonary hypertension by a pulsed Doppler technique. Circulation 1983; 68(2): 302-309.