

# Calculation of the vector modes of an optical waveguide

V.V.Kotlyar<sup>1,2</sup>, Y.O.Shuyupova<sup>2</sup>

<sup>1</sup>*Image Processing Systems Institute of RAS*

<sup>2</sup>*Samara State Aerospace University (SSAU)*

## **Abstract:**

The paper considers a method for calculation of eigenmodes of waveguides based on the method of conjugated sinusoidal modes. The corresponding software developed in the MathLab environment is used for the calculation of the modes of a weakly direct step-index fiber.

**Keywords:** optical waveguide, conjugated sinusoidal modes, MathLab, weakly direct step-index

**Acknowledgments:** This work was carried out as part of the Russian-American program “Basic Research and Higher Education” (BRHE), and was also supported by the Presidential grant of the Russian Federation NSh-1007.2003.01.

**Citation:** Kotlyar VV, Shuyupova YO. Calculation of the vector modes of an optical waveguide. Computer Optics 2005; 27: 89-94.

[Access full text \(in Russian\)](#)

## **References:**

- [1] Kotlyar VV, Shuyupova YO. Calculation of the spatial optical waveguide modes with inhomogeneous cross-sections using the sinusoidal mode matching method. Computer Optics 2003; 25: 41-48.
- [2] Sudbo ASv. Film mode matching: a versatile method for mode film calculations in dielectric waveguides. Pure Appl Opt 1993; 2(3): 211-233 DOI: 10.1088/0963-9659/2/3/007..
- [3] Sudbo ASv. Improved formulation of the film mode matching method for mode film calculations in dielectric waveguides. Pure Appl Opt 1994; 3(3): 381-388. DOI: 10.1088/0963-9659/3/3/021.
- [4] Stagira S. Full vectorial analysis of cylindrical waveguides using Green functions. Opt Commun 2003; 225(4-6): 281-291. DOI: 10.1016/j.optcom.2003.08.009.
- [5] Lusse P, Stuwe P, Schule J, Unger H. Analysis of vectorial mode fields in optical waveguides by a new finite difference method. J Light Technol 1994; 12(3): 487-494. DOI: 10.1109/50.285331.
- [6] Hadley GR, Smith RE. Full-vector waveguide modeling using an iterative finite-difference method with transparent boundary conditions. J Light Technol 1994; 12(3): 487-494. DOI: 10.1109/50.285331.
- [7] Rogge U, Pregla R. Method of lines for the analysis of dielectric waveguides. J Light Technol 1993; 11(12): 2015-2020. DOI: 10.1109/50.257964.
- [8] Sztefka G, Nogling HP. Bidirectional eigenmode propagation for large refractive index steps. IEEE Photonics Technol Lett 1993; 5(5): 554-557. DOI: 10.1109/68.215279.
- [9] Rahman BMA, Davies JB. Finite-Elements Solution of integrated optical waveguides. J Light Technol 1984; 2(5): 682-687. DOI: 10.1109/JLT.1984.1073669.
- [10] Koshiba M, Maruyama S, Hirayama K. A vector finite element method with the high-order mixedinterpolation-type triangular elements for optical waveguiding problems. J Light Technol 1994; 12(3): 495-502. DOI: 10.1109/50.285332.
- [11] Lusse P, Stuwe P, Schule J, Unger H-G. Analysis of vectorial mode fields in optical waveguides by a new finite difference method. J Light Technol 1994; 12(3): 487-494. DOI: 10.1109/50.285331.
- [12] Hadley GR, Smith RE. Full-vector waveguide modeling using an iterative finite-difference method with transparent boundary conditions. J Light Technol 1995; 13(3): 465-469. DOI: 10.1109/50.372444.
- [13] Lin P-L, Li B-J. Semivectorial Helmholtz beam propagations by Lanczos reduction. IEEE J Quantum Electron 1993; 29(8): 2385-2389. DOI: 10.1109/3.245570.
- [14] Lee P-C, Voges E. Three dimensional semi-vectorial wide-angle beam propagation method. J Light Technol 1994; 12(2): 215-225. DOI: 10.1109/50.350601.